

# A VALUATION CRITERION FOR NORMAL BASIS GENERATORS IN LOCAL FIELDS OF CHARACTERISTIC $p$

G. GRIFFITH ELDER

**ABSTRACT.** Let  $K$  be a complete local field of characteristic  $p$  with perfect residue field. Let  $L/K$  be a finite, fully ramified, Galois  $p$ -extension. If  $\pi_L \in L$  is a prime element, and  $p'(x)$  is the derivative of  $\pi_L$ 's minimal polynomial over  $K$ , then the relative different  $\mathfrak{D}_{L/K}$  is generated by  $p'(\pi_L) \in L$ . Let  $v_L$  be the normalized valuation normalized with  $v_L(L) = \mathbb{Z}$ . We show that any element  $\rho \in L$  with  $v_L(\rho) \equiv -v_L(p'(\pi_L)) - 1 \pmod{[L : K]}$  generates a normal basis,  $K[\text{Gal}(L/K)] \cdot \rho = L$ . This criterion is tight: Given any integer  $i$  such that  $i \not\equiv -v_L(p'(\pi_L)) - 1 \pmod{[L : K]}$ , there is a  $\rho_i \in L$  with  $v_L(\rho_i) = i$  such that  $K[\text{Gal}(L/K)] \cdot \rho_i \subsetneq L$ .

The Normal Basis Theorem states that in a finite Galois extension  $L/K$  with  $G = \text{Gal}(L/K)$ , there is an element  $\rho \in L$ , called a normal basis generator, whose conjugates  $\{\sigma\rho : \sigma \in G\}$  provide a basis for  $L$  over  $K$ . In the setting of local field extensions, the most important property of an element is its valuation, and so [2] asked whether there is a valuation criterion: Is there a valuation (an integer certificate) that guarantees that any element bearing this valuation is a normal basis generator?

Let  $K$  be a complete local field with a perfect residue field of characteristic  $p$ . So the characteristic of  $K$  is 0 or  $p$ . It is not too hard to see that as a necessary condition, if a valuation criterion exists for a finite Galois extension  $L/K$ , then  $L/K$  must be fully ramified and have order a power of  $p$ . But are these necessary conditions also sufficient? In this paper we give an affirmative answer to this question in the case where  $K$  has characteristic  $p$ .

For  $K$  of characteristic 0 and regular (so  $K$  does not contain the  $p$ th roots of unity), a valuation criterion for fully ramified elementary abelian  $p$ -extensions is given in [2]. For  $K$  of characteristic  $p$ , a valuation criterion for fully ramified abelian  $p$ -extensions is given in [5]. In both cases, the valuation criterion is described in terms of the largest ramification break number associated with  $L/K$ . The main contribution of this paper is a restatement of that criterion in terms of the exponent of the relative different (see remark following proposition). The relative different satisfies  $\mathfrak{D}_{L/K} = (p'(\pi_L))$  with  $\pi_L$  a prime element of  $L$  and  $p(x)$  the minimal polynomial of  $\pi_L$  over  $K$  [4, III §6 Lemma 2 Corollary 2]. This means that we can state our main result as follows:

**Theorem 0.1.** *Let  $K$  be a complete local field of characteristic  $p$  with perfect residue field. Let  $L/K$  be a finite, Galois extension with  $G = \text{Gal}(L/K)$ , and let  $v_L$  be the valuation normalized so that  $v_L(L) = \mathbb{Z}$ . If  $L/K$  is fully ramified and  $[L : K] = p^n$  for some integer  $n$ , there is a valuation criterion: Let  $\pi_L \in L$  be a*

---

*Date:* February 12, 2008.

*1991 Mathematics Subject Classification.* 11S15.

prime element and let  $p(x)$  be its minimal polynomial over  $K$ . Then any element  $\rho \in L$  with  $v_L(\rho) \equiv -v_L(p'(\pi_L)) - 1 \pmod{[L : K]}$  generates a normal basis for  $L/K$ . So  $L = K[G] \cdot \rho$ .

Moreover the assumption that  $L/K$  is a fully ramified  $p$ -extension is necessary and the criterion under that assumption is tight: Outside of the assumption, given any  $i \in \mathbb{Z}$ , or under the assumption, given any  $i \not\equiv -v_L(p'(\pi_L)) - 1 \pmod{[L : K]}$ , there is a  $\rho_i \in L$  with  $v_L(\rho_i) = i$  such that  $L \supsetneq K[G] \cdot \rho_i$ .

*Proof.* Recall the definition of the ramification groups  $G_i$  [4, IV]. We begin by assuming that  $G = G_1$ , which is equivalent to the assumption that  $L/K$  is a fully ramified extension of degree  $p^n$  for some  $n$ . Let  $d = v_L(p'(\pi_L))$  and let  $\rho \in L$  with  $v_L(\rho) \equiv -d - 1 \pmod{p^n}$ . Express  $\rho$  in terms of the field basis  $B = \{\pi_L^i/p'(\pi_L) : i = 0, \dots, p^n - 1\}$  for  $L/K$ . It is a result of Euler that

$$\mathrm{Tr}_G \frac{\pi_L^i}{p'(\pi_L)} = \begin{cases} 0 & \text{for } 0 \leq i \leq p^n - 2 \\ 1 & \text{for } i = p^n - 1 \end{cases}$$

[4, III §6 Lemma 2]. Since  $v_L(\rho) \equiv v_L(\pi_L^{p^n-1}/p'(\pi_L)) \pmod{p^n}$ , the coefficient of  $\pi_L^{p^n-1}/p'(\pi_L)$  in the expression for  $\rho$  must be nonzero. Therefore  $\mathrm{Tr}_G \rho \neq 0$ . Moreover, we can replace  $B$  by the alternate basis  $\{\rho, \pi_L^i/p'(\pi_L) : i = 0, \dots, p^n - 2\}$ , and get

$$L = K \cdot \rho + \sum_{i=0}^{p^n-2} K \cdot \frac{\pi_L^i}{p'(\pi_L)}.$$

The Normal Basis Theorem, stated in terms of Tate cohomology, says that  $\hat{H}^{-1}(G, L) = 0$  [4, VIII §1 & X §1 Proposition 1]. This means that any element  $\eta \in L$  with  $\mathrm{Tr}_G \eta = 0$  satisfies  $\eta \in \mathcal{I}_G \cdot L$ , where  $\mathcal{I}_G = (\sigma - 1 : \sigma \in G)$  is the augmentation ideal of the group ring  $K[G]$ . Therefore  $L = K \cdot \rho + \mathcal{I}_G \cdot L$ , but also

$$L = K[G] \cdot \rho + \mathcal{I}_G \cdot L.$$

Now notice that because  $G$  is a  $p$ -group and  $K$  has characteristic  $p$ ,  $\mathcal{I}_G$  is also the Jacobson radical of  $K[G]$ . Thus, by Nakayama's Lemma,  $L = K[G] \cdot \rho$ . We have proven the criterion. The elements  $\pi_K^m \pi_L^{p^n-1}/p'(\pi_L)$  for  $i \neq p^n - 1$  show that the criterion is sharp. (All of this argument applies equally well in characteristic 0, except for one step: In characteristic 0, the augmentation ideal  $\mathcal{I}_G$  is not the Jacobson radical of  $K[G]$ .)

Now we address  $G \supsetneq G_1$ . Let  $L' = L^{G_1}$  and  $L'' = L^{G_0}$ . So  $L''/K$  is unramified and  $L'/L''$  is fully but tamely ramified. Replace  $K$  and  $G$ , in our earlier argument with  $L'$  and  $G_1$  respectively. So  $d$  is defined in terms of the relative different of the fully ramified  $p$ -part of the extension,  $\mathfrak{P}_L^d = \mathfrak{D}_{L/L'}$  and  $p^n = [L : L']$ . From earlier work, given any integer  $i \not\equiv -d - 1 \pmod{p^n}$  there is a  $\rho \in L$  with  $v_L(\rho) = i$  such that  $\mathrm{Tr}_{G_1} \rho = 0$ . So  $K[G] \rho \subsetneq L$ .

To consider the case  $i \equiv -d - 1 \pmod{p^n}$ , notice that the trace  $\mathrm{Tr}_{G_1}$  maps fractional ideals of  $\mathfrak{D}_L$ , the ring of integers of  $L$ , to fractional ideals of  $\mathfrak{D}_{L''}$ . Indeed, using basis  $B$  from above,  $\mathrm{Tr}_{G_1} \mathfrak{P}_L^{kp^n-d-1} = \mathfrak{P}_{L'}^{k-1}$ . Moreover  $\mathrm{Tr}_{G_1} \mathfrak{P}_L^{kp^n-d} \subseteq \mathfrak{P}_{L'}^k$ , because of [4, III §3 Proposition 7]. Observe that this means that given any element  $\tau \in L'$  with  $v_{L'}(\tau) = k - 1$ , there is a  $\rho_\tau \in L$  with  $v_L(\rho_\tau) = kp^n - d - 1$  such that  $\mathrm{Tr}_{G_1}(\rho_\tau) = \tau$ .

To use this observation notice that because  $L'/L''$  is tamely ramified, there is a prime element of  $L''$ , namely  $\pi_{L''}$ , such that  $L' = L''(\sqrt[p]{\pi_{L''}})$  with  $p \nmid e$  [3, II §3.5

Proposition]. So for  $k \not\equiv 1 \pmod{e}$ , let  $\tau = \sqrt[e]{\pi_{L''}}^{k-1}$ . Since  $\text{Tr}_{G_0/G_1} \sqrt[e]{\pi_{L''}}^{k-1} = 0$ ,  $\text{Tr}_{G_0} \rho = 0$  and  $K[G]\rho \subsetneq L$ . For  $k \equiv 1 \pmod{e}$ , let  $\tau = \pi_K^{(k-1)/e}$  and let  $\bar{\sigma}$  be any non-trivial element in  $G/G_1$ . Then  $(\sigma - 1)\pi_K^{k-1} = 0$ . Thus  $(\sigma - 1)\text{Tr}_{G_1} \rho = 0$  and  $K[G]\rho \subsetneq L$ .  $\square$

Now we connect the valuation criterion of this paper with that of [2], [5].

**Proposition 0.2.** *Let  $K$  be a complete local field (characteristic 0 or  $p$ ) with perfect residue field of characteristic  $p$ . Let  $L/K$  be a finite, fully ramified, Galois  $p$ -extension. Let  $\pi_L \in L$  be a prime element and let  $p(x)$  be its minimal polynomial over  $K$ . Then*

$$v_L(p'(\pi_L)) + 1 \equiv p^n u_m - b_m \pmod{[L : K]}$$

where  $b_m, u_m$  are the largest ramification break numbers in lower and upper numbering respectively.

**Remark 1.** When  $L/K$  is abelian  $u_m \in \mathbb{Z}$  by the Hasse-Arf Theorem and so we get the valuation criterion of [2], [5], namely  $v_L(\rho) \equiv b_m \pmod{p^n}$ .

*Proof.* Let  $d = v_L(p'(\pi_L))$  and recall that  $\mathfrak{D}_{L/K} = (p'(\pi_L))$ . Let  $G = G_1$  and  $[L : K] = p^n$ . Recall the ramification filtration  $G = G_1 \supseteq G_2 \supseteq \cdots$  where  $G_i = \{\sigma \in G : v_L((\sigma - 1)\pi_L) \geq i + 1\}$  [4, IV §1]. The break numbers (in lower numbering) are those integers  $i$  such that  $G_i \supsetneq G_{i+1}$ . Let  $b_1 < b_2 < \cdots < b_m$  be the list of break numbers with  $b_m$  being the maximal break (again in lower numbering). Let  $g_i = |G_{b_i}|$ , the number of elements in  $G_{b_i}$ . Then by [4, IV §1 Proposition 4],

$$d = (1 + b_1)(g_1 - 1) + \sum_{i=2}^m (b_i - b_{i-1})(g_i - 1) = (1 + b_1)g_1 - b_m - 1 + \sum_{i=2}^m (b_i - b_{i-1})g_i.$$

Moreover, we can convert the lower numbering to upper numbering using the Herbrand function  $\varphi$  [4, IV §3]. Be careful to notice a small difference in notation: We use  $g_i = |G_{b_i}|$  whereas [4, IV §3] uses  $g_{b_i} = |G_{b_i}|$ . The largest break number in upper numbering is therefore

$$u_m = \varphi(b_m) = \frac{1}{p^n} \left( b_1 g_1 + \sum_{i=2}^m (b_i - b_{i-1}) g_i \right).$$

Thus  $d + 1 = g_1 - b_m + p^n u_m$ , where  $g_1 = p^n$ .  $\square$

We end the paper with a natural

**Question 1.** Does the statement of the Theorem (modified appropriately to address [2, Example 1]) also hold in characteristic zero?

## REFERENCES

- [1] B. de Smit and L. Thomas, ‘Local Galois module structure in positive characteristic and continued fractions’, *Arch. Math. (Basel)* 88 (2007), no. 3, 207–219.
- [2] N. P. Byott and G. G. Elder, ‘A valuation criterion for normal bases in elementary abelian extensions’, *Bull. Lond. Math. Soc.* 39 (2007), no. 5, 705–708.
- [3] I. B. Fesenko and S. V. Vostokov, ‘Local fields and their extensions’, (American Mathematical Society, Providence RI, 2002).
- [4] J.-P. Serre, ‘Local fields’, (Springer-Verlag, New York, 1979).
- [5] L. Thomas, ‘A valuation criterion for normal basis generators in equal positive characteristic’, preprint: August 9, 2007.

*E-mail address:* `elder@unomaha.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEBRASKA AT OMAHA, OMAHA, NE 68182-0243 U.S.A.